



# COMMENTS ON THE ARTICLE: "AGENTS RELATED TO THE SURGICAL SITE INFECTION OF POSTOPERATIVE ELDERLY PEOPLE AT EL CENTRO MEDICO NAVAL" 2013-2017"

COMENTARIOS SOBRE EL ARTÍCULO: "AGENTES RELACIONADOS A INFECCIÓN DE SITIO OPERATORIO EN ADULTOS MAYORES POS OPERADOS EN EL CENTRO MÉDICO NAVAL, 2013 – 2017"

Jordy Pulido-Flores<sup>1</sup>, Andre Ore-Arce<sup>1</sup>, Jacqueline Reyes-Gamonal<sup>1</sup>

## Mr. Editor

After reading this article with interest published by Hidalgo et al<sup>1</sup> regarding agents related to the surgical site infection (SSI) in elderly between the years 2013 and 2017; we found the consideration of certain relevant determinants related to clinical and surgical aspects. These determinants are useful since they are taken as reference in future research work. Nevertheless, we would like to emphasize the necessity to incorporate some other relevant variables as possible determinants to control such as those related to nutritional status specifically hemoglobin and preoperative albumin levels since in our country this kind of pathologies is common.

In different countries<sup>2</sup>, SSI is one of the most deadly and costly complications increasing mortality from 2 to 11 times<sup>3</sup> and the cost to 14.2 million dollars in surgeries<sup>4</sup>. In Peru, from 15 000 nosocomial infections, 30, 9 % are from an surgical wound<sup>5</sup>. It is important to know the details of this theme because it contributes to a highly reversible social and economic problem.

The nutritional status of the elderly is vital to keep an adequate immune response<sup>6</sup>. A common alteration is anemia, a frequent public health problem in underdeveloped countries such as Peru<sup>7</sup>. It constitutes a significant risk factor that can increase the person fragility<sup>8,9</sup> and thus increase a SSI development likelihood<sup>10</sup>. As evidenced by the study of Atalaya Marín in which from 49.5% of anemic patients, 67.9% present post-surgical complications in which the SSI stands out<sup>11</sup>.

On the other hand, albumin values are in general very useful for determining the patient's nutritional status as it is available in hospitals<sup>12</sup>. Poor albumin levels have been related to patients with scarring alterations due to altered collagen synthesis and thus it prevents granuloma formation in surgical wounds predisposing to bacterial contamination in the hospital environment<sup>13</sup>. Although it is known that albumin is reduced in acute phase response, conclusions cannot be drawn as to whether preoperative prealbumin levels could predict an infection development<sup>14</sup>. In spite of this, different studies have described that albumin deficiency increases the SSI risk up to 12.6 times more<sup>13</sup>. Furthermore, it is stated that patients with postoperative serum albumin values <3.5 g / dL have a significantly higher risk of SSI mortality.

Even though the Peruvian state is dealing with malnutrition, it is necessary to intensify efforts to prevent these types of complications, which are socially and economically damaging. We believe it is necessary to emphasize the person's nutritional status as a risk factor to develop SSI in future work to be developed.

<sup>1</sup> Faculty of Human Medicine, Ricardo Palma University, Lima-Peru.

Cite as: Jordy Pulido-Flores, Andre Ore-Arce, Jacqueline Reyes-Gamonal. Comments on the article: "Agents related to the surgical site infection of postoperative elderly people at El Centro Medico Naval" 2013-2017". Rev. Fac. Med. Hum. October 2019; 19(4):138-139. DOI 10.25176/RFMH.v19i4.2348

**Authorship Contributions:** The authors participated in the generation, collection of information, writing and final version of the original article.

**Financing:** Self-financed

**Interest conflict:** The authors declare no conflicts of interest in the publication of this article.

**Received:** August 28, 2019

**Approved:** September 10, 2019

**Correspondence:** Jordy Pulido-Flores

**Address:** Av. Alfredo Benavides 5440, Santiago de Surco, Lima-Perú.

**Telephone:** 920238351

**E-mail:** jpulidoflores@gmail.com

## BIBLIOGRAPHIC REFERENCES

1. Lucía Fernanda Hidalgo-Vizcarreta, Magdiel José Manuel Gonzales-Menéndez, Cecilia Roxana Salinas-Salas. Agentes relacionados a infección de sitio operatorio en adultos mayores pos operados en el centro médico naval, 2013 – 2017. Rev. Fac. Med. Hum. Julio 2019; 19(3):43-47. Disponible en: 10.25176/RFMH.v19i3.2163
2. Berrios-Torres SI, Umscheid CA, Bratzler DW, Leas B, Stone EC, Kelz RR, et al. Centers for Disease Control and Prevention Guideline for the Prevention of Surgical Site Infection, 2017. JAMA Surg. 1 de agosto de 2017;152(8):784-91. Disponible en: 10.1001/jamasurg.2017.0904
3. Ban KA, Minei JP, Laronga C, Harbrecht BG, Jensen EH, Fry DE, et al. American College of Surgeons and Surgical Infection Society: Surgical Site Infection Guidelines, 2016 Update. Journal of the American College of Surgeons. 1 de enero de 2017;224(1):59-74. Disponible en: 10.1016/j.jamcollsurg.2016.10.029
4. National Healthcare Safety Network, Centers for Disease Control and Prevention. Surgical site infection (SSI)event. January 2017. Accessed January 25, 2017. Disponible en: <https://www.cdc.gov/nhsn/PDFs/pscManual/9pscSSIconcurrent.pdf>
5. Bonifacio M, María S. Malnutrición como factor de riesgo asociado a las infecciones de sitio operatorio del departamento de Cirugía general del Hospital Nacional Daniel Alcides Carrión en el periodo enero-junio 2017. Universidad Ricardo Palma [Internet]. 2018 [citado 11 de septiembre de 2019]; Disponible en: <http://repositorio.urp.edu.pe/handle/URP/1178>
6. López Plaza B, Bermejo López LM. Nutrición y trastornos del sistema inmune. Nutrición Hospitalaria. 2017;34:68-71. Disponible en: <http://dx.doi.org/10.20960/nh.1575>.
7. Tarqui-Mamani C, Sanchez-Abanto J, Alvarez-Dongo D, Espinoza-Oriundo P, Jordan-Lechuga T. Prevalencia de anemia y factores asociados en adultos mayores peruanos. Rev Peru Med Exp Salud Publica. diciembre de 2015;32:687-92. Disponible en: [https://www.scielosp.org/scielo.php?pid=S1726-46342015000400009&script=sci\\_arttext&tlang=en](https://www.scielosp.org/scielo.php?pid=S1726-46342015000400009&script=sci_arttext&tlang=en)
8. Stauder R, Valent P, Theurl I. Anemia at older age: etiologies, clinical implications, and management. Blood. 1 de febrero de 2018;131(5):505-14. Disponible en: <https://doi.org/10.1182/blood-2017-07-746446>
9. Pires Corona L, Drumond Andrade FC, de Oliveira Duarte YA, Lebrao ML. The relationship between anemia, hemoglobin concentration and frailty in Brazilian older adults. J Nutr Health Aging. 1 de noviembre de 2015;19(9):935-40. Disponible en: <https://doi.org/10.1007/s12603-015-0502-3>
10. Sattar F, Sattar Z, Zaman M, Akbar S. Frequency of Post-operative Surgical Site Infections in a Tertiary Care Hospital in Abbottabad, Pakistan. Cureus. 2019;11(3). DISPONIBLE EN: 10.7759/cureus.4243
11. Marin A, Roosevelt E. El hemograma completo como parámetro predictor de riesgo de infección en sitio operatorio, en los pacientes operados en el servicio de cirugía del HRDC en el año 2017. Universidad Nacional de Cajamarca. 2019. Disponible en: <http://repositorio.unc.edu.pe/handle/UNC/2963>
12. Molina Soria JB, Lobo Támer G, Pérez de la Cruz AJ, Ruiz-López MD. Prevalencia de desnutrición al ingreso en un hospital general básico. Nutr Hosp 2017;34:1390-1398. Disponible en: <http://dx.doi.org/10.20960/nh.1133>
13. Bacardí Zapata PA, Páez Candelaria Y, Jones Romero O, Romero García LI, Ricardo Ramírez JM, Gondres Legró KM. Hipoalbuminemia e infección postoperatoria en una unidad de atención al grave. Cuba y Salud. 2017;12(3):24-9.n Disponible en : <https://www.medicgraphic.com/pdfs/cubaysalud/pcs-2017/pcs173e.pdf>
14. Salvetti DJ, Tempel ZJ, Gandhoke GS, Parry PV, Grandhi RM, Kanter AS, et al. Preoperative prealbumin level as a risk factor for surgical site infection following elective spine surgery. Surg Neurol Int. 8 de octubre de 2015;6(Suppl 19):S500-3. DISPONIBLE en: 10.4103/2152-7806.166893